

FIBROSIS

iRhom2 and TNF: Partners or enemies?

Marina Badenes^{1*} and Colin Adrain^{1,2*}

iRhom2 is an essential cofactor for ADAM17, the metalloprotease that sheds both the proinflammatory cytokine tumor necrosis factor- α (TNF- α) and TNF receptors (TNFRs) from the cell surface. In this issue of *Science Signaling*, Sundaram *et al.* demonstrate a protective role for iRhom2 in promoting ADAM17-mediated shedding of TNFRs in hepatic stellate cells, which reduces TNFR signaling and liver fibrosis in response to injury.

Copyright © 2019
The Authors, some
rights reserved;
exclusive licensee
American Association
for the Advancement
of Science. No claim
to original U.S.
Government Works

ADAM17 [A disintegrin and metalloprotease 17, also called tumor necrosis factor- α (TNF- α)-converting enzyme (TACE)] is a pleiotropic protease that cleaves many membrane-tethered substrates, including the proinflammatory cytokine TNF- α and its receptors TNFR1 and TNFR2 (1). The pseudo-protease iRhom2 is an essential cofactor for ADAM17 that is highly abundant in immune cells (2). iRhom2 is required for multiple aspects of ADAM17 biology, including the trafficking of ADAM17 into the Golgi apparatus, where it undergoes a key proteolytic maturation step, the stimulation of ADAM17 activity on the cell surface (3), and the control of ADAM17 proteolytic specificity (4). iRhom2 deficiency protects mice from several inflammatory diseases by blocking ADAM17-mediated TNF- α secretion in inflammatory cells (2, 5). The current assumption in the field, therefore, is that iRhom2 is proinflammatory. In this issue, Sundaram and co-workers (6) demonstrate that iRhom2 plays a protective role in a mouse model of inflammatory liver injury and fibrosis by attenuating TNFR signaling.

Liver fibrosis develops as a result of chronic liver injury and can lead to liver cirrhosis and hepatocellular carcinoma. During liver injury, the release of inflammatory mediators, such as TNF- α and interleukin-1 (IL-1), activates nuclear factor κ B (NF- κ B) to promote the proliferation of a normally quiescent population of cells called hepatic stellate cells (HSCs) (7). This triggers the proliferation and differentiation of HSCs into profibrotic myofibroblasts that secrete collagen, thus promoting liver fibrosis (8).

The authors of the present study found that human patients with liver cirrhosis and mice with liver fibrosis induced by bile duct ligation (BDL) both showed increased circulating amounts of TNFR1 and TNFR2 and

increased expression of *Rhbd2*, the gene that encodes iRhom2, in the liver. Ex vivo experiments with murine HSCs showed that iRhom2 was required for the shedding of both TNFR1 and TNFR2. Consistent with this, in mice subjected to BDL, deletion of *Rhbd2* increased TNF- α signaling in the liver, presumably by blocking the shedding of TNFRs. This, in turn, was associated with HSC proliferation and fibrosis potentially mediated by augmented TNFR1 signaling. Consistent with this hypothesis, treatment of *Rhbd2* knockout mice with the TNF- α inhibitor etanercept blocked HSC proliferation and ameliorated liver fibrosis. The authors conclude that following BDL, the increased expression of *Rhbd2* in the liver augmented ADAM17 activity, leading to the shedding of TNFRs, which dampened TNF- α signaling and prevented the proliferation of HSCs and the development of liver fibrosis (Fig. 1). In contrast to the conventional view of iRhom2 as proinflammatory, these findings reveal a scenario in which iRhom2 attenuates signaling through an inflammatory signaling pathway to limit overactivation of tissue repair pathways that could lead to fibrosis.

Because TNFR1 and TNFR2 can exert distinct biological outcomes, an obvious question concerns which TNFR is responsible for promoting the NF- κ B-dependent signaling pathways that make HSCs profibrogenic. Although Sundaram and colleagues (6) showed that HSCs release both TNFR1 and TNFR2 in an iRhom2-dependent manner, when stimulated with TNF- α to mimic an inflammatory environment, TNFR2 was preferentially cleaved. This suggests a model whereby TNFR2 is shed as a decoy receptor, preventing TNF- α signaling through TNFR1. This is the same mechanism by which etanercept, an engineered soluble form of TNFR2, acts. These data are consistent with previous studies

that demonstrated a requirement for TNFR1, but not TNFR2, in protection from BDL-triggered fibrosis (7). Notably, TNFR1 activation can promote fibrogenesis by inducing the remodeling of the extracellular matrix by stimulating the production of matrix metalloproteinase-9 (MMP9) through activation of the NF- κ B pathway (7). Therefore, the profibrotic effects of iRhom2 deletion observed in the current paper can likely be attributed to TNFR1 signaling.

What this work highlights more generally is that iRhoms and ADAM17 can impinge on TNF- α signaling in multiple ways, leading to sophisticated biological scenarios. First, the shedding of soluble TNF- α (sTNF- α), which is proinflammatory and preferentially activates TNFR1, could drive inflammation—the widely reported role of iRhom2. Second, the shedding of TNFR1 and/or TNFR2 could attenuate TNF- α signaling by blocking signal reception in cells with reduced membrane TNFRs, thus reducing signal flux. Alternatively, sTNFRs could act as a diffusible sink for sTNF- α , blocking signaling in a non-cell-autonomous manner. Finally, the failure to produce sTNF- α would result in the accumulation of membrane-tethered TNF- α (mTNF- α), which acts preferentially on TNFR2 (9), eliciting distinct biological outcomes from those elicited by TNFR1 activation (10). In the latter case, iRhoms, rather than tuning signaling flux, could mediate a form of mode switching in the nature of TNF- α signaling.

Added to this already complicated scenario is the fact that mammals have a second iRhom paralog, iRhom1, with a distinct but partially overlapping tissue distribution and a shared function with iRhom2 in ADAM17 regulation. Furthermore, iRhom1 and iRhom2 can differentially govern ADAM17 substrate selectivity (4), providing another potential basis by which iRhoms can qualify the nature of TNF- α signaling by promoting the cleavage of a specific TNFR preferentially in a given tissue. Finally, iRhoms are regulated

¹Instituto Gulbenkian de Ciência (IGC), Oeiras, Portugal. ²Centre for Cancer Research and Cell Biology, Queen's University, Belfast, UK.

*Corresponding author. Email: mbadenes@igc.gulbenkian.pt (M.B.); cadrain@igc.gulbenkian.pt (C.A.)




Fig. 1. Mechanism by which iRhom2 attenuates HSC activation and fibrogenesis. During liver injury, Kupffer cells, monocyte-derived macrophages, or hepatocytes secrete inflammatory cytokines, including TNF- α . The resulting inflammatory environment leads to an increase in the amounts of iRhom2 in hepatic stellate cells (HSCs), increasing ADAM17 activity and leading to the cleavage of TNFR2. This allows the latter to act as a soluble decoy to sequester TNF- α , thus preventing TNFR1 signaling. In the absence of this iRhom2-dependent inhibitory mechanism, TNF- α binds to TNFR1 to promote HSC proliferation and activation to become myofibroblasts, leading to fibrosis.

at both the transcriptional and posttranscriptional levels, providing a sophisticated manner in which, depending on the stimulus or developmental context, the amounts of an individual iRhom in a given tissue may be

modulated to control signaling in a subtle—or indeed fundamental—manner.

A full understanding of how iRhom2 regulates TNF- α signaling has been a challenge for the above-noted reasons. Comparing the iRhom2 knockout mice with TNF- α , TNFR1 and TNFR2 mutants may not give sufficient clarity because of the complexity in the iRhom regulation of TNF- α signaling. In the case of the present paper, the deletion of iRhom2 or ADAM17 specifically in HSCs would clarify if, in BDL-induced liver fibrosis, the protective effect of iRhom2 is mediated by its action in HSCs or whether other relevant cell types that also produce iRhom2 (such as hepatocytes, Kupffer cells, or monocyte-derived macrophages) are involved. It would also be interesting to determine whether iRhom2 exerts a protective role in different models of liver fibrosis and also in other TNF- α -mediated diseases. Nevertheless, this work reinforces the notion that iRhoms, in conjunction with ADAM17, can modulate TNF- α signaling in a highly sophisticated manner that can lead to signal attenuation, potentiation, or mode switching.

REFERENCES AND NOTES

- J. Scheller, A. Chalaris, C. Garbers, S. Rose-John, ADAM17: A molecular switch to control inflammation and tissue regeneration. *Trends Immunol.* **32**, 380–387 (2011).
- C. Adrain, M. Zettl, Y. Christova, N. Taylor, M. Freeman, Tumor necrosis factor signaling requires iRhom2 to promote trafficking and activation of TACE. *Science* **335**, 225–228 (2012).
- M. Cavadas, I. Oikonomidi, C. J. Gaspar, E. Burbridge, M. Badenes, I. Felix, A. Bolado, T. Hu, A. Bileck, C. Gerner, P. M. Domingos, A. von Kriegsheim, C. Adrain, Phosphorylation of iRhom2 controls stimulated proteolytic shedding by the metalloprotease ADAM17/TACE. *Cell Rep.* **21**, 745–757 (2017).
- T. Maretzky, D. R. McIlwain, P. D. Issuree, X. Li, J. Malapeira, S. Amin, P. A. Lang, T. W. Mak, C. P. Blobel, iRhom2 controls the substrate selectivity of stimulated ADAM17-dependent ectodomain shedding. *Proc. Natl. Acad. Sci. U.S.A.* **110**, 11433–11438 (2013).
- P. D. A. Issuree, T. Maretzky, D. R. McIlwain, S. Monette, X. P. Qing, P. A. Lang, S. L. Swendeman, K. H. Park-Min, N. Binder, G. D. Kalliolias, A. Yarinina, K. Horiuchi, L. B. Ivashkiv, T. W. Mak, J. E. Salmon, C. P. Blobel, iRHOM2 is a critical pathogenic mediator of inflammatory arthritis. *J. Clin. Invest.* **123**, 928–932 (2013).
- B. Sundaram, K. Behnke, A. Belancic, M. A. Al-Salihi, Y. Thabet, R. Polz, R. Pellegrino, Y. Zhuang, P. V. Shinde, H. C. Xu, J. Vasilevska, T. Longenich, D. Herebian, E. Mayatepek, H. H. Bock, P. May, C. Kordes, N. Aghaeepour, T. W. Mak, V. Keitel, D. Häussinger, J. Scheller, A. A. Pandya, K. S. Lang, P. A. Lang, iRhom2 prevents bile duct obstruction-induced liver fibrosis. *Sci. Signal.* **12**, eaax1194 (2019).
- N. Tarrats, A. Moles, A. Morales, C. Garcia-Ruiz, J. C. Fernandez-Checa, M. Mari, Critical role of tumor necrosis factor receptor 1, but not 2, in hepatic stellate cell proliferation, extracellular matrix remodeling, and liver fibrogenesis. *Hepatology* **54**, 319–327 (2011).
- C.-Y. Zhang, W.-G. Yuan, P. He, J.-H. Lei, C.-X. Wang, Liver fibrosis and hepatic stellate cells: Etiology, pathological hallmarks and therapeutic targets. *World J. Gastroenterol.* **22**, 10512–10522 (2016).
- M. Grell, E. Douni, H. Wajant, M. Lohden, M. Claus, B. Maxeiner, S. Georgopoulos, W. Lesslauer, G. Kollias, K. Pfizenmaier, P. Scheurich, The transmembrane form of tumor necrosis factor is the prime activating ligand of the 80 kDa tumor necrosis factor receptor. *Cell* **83**, 793–802 (1995).
- T. Hehlhans, K. Pfeffer, The intriguing biology of the tumour necrosis factor/tumour necrosis factor receptor superfamily: Players, rules and the games. *Immunology* **115**, 1–20 (2005).

Acknowledgments: We thank the members of our laboratory for stimulating discussions. **Funding:** We are grateful for the ongoing support of our laboratory by Fundação Calouste Gulbenkian, Queen's University Belfast, and Fundação para a Ciência e Tecnologia (project LISBOA-01-0145-FEDER-031330). The project leading to these results has received funding from “la Caixa” Foundation (ID 100010434), under the agreement <LCF/PR/HR17/52150018>.

Competing interests: The authors declare that they have no competing interests.

10.1126/scisignal.aaz0444

Citation: M. Badenes, C. Adrain, iRhom2 and TNF: Partners or enemies? *Sci. Signal.* **12**, eaaz0444 (2019).

iRhom2 and TNF: Partners or enemies?

Marina Badenes and Colin Adrain

Sci. Signal. **12** (605), eaaz0444.
DOI: 10.1126/scisignal.aaz0444

ARTICLE TOOLS

<http://stke.sciencemag.org/content/12/605/eaaz0444>

RELATED CONTENT

<http://stke.sciencemag.org/content/sigtrans/12/564/eaao3469.full>
<http://stke.sciencemag.org/content/sigtrans/11/511/eaao4910.full>
<http://stke.sciencemag.org/content/sigtrans/12/592/eaaw2418.full>
<http://stke.sciencemag.org/content/sigtrans/12/605/eaax1194.full>
<http://immunology.sciencemag.org/content/immunology/3/28/eaar7754.full>

REFERENCES

This article cites 10 articles, 3 of which you can access for free
<http://stke.sciencemag.org/content/12/605/eaaz0444#BIBL>

PERMISSIONS

<http://www.sciencemag.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of Service](#)

Science Signaling (ISSN 1937-9145) is published by the American Association for the Advancement of Science, 1200 New York Avenue NW, Washington, DC 20005. The title *Science Signaling* is a registered trademark of AAAS.

Copyright © 2019 The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works