

Sleep deprivation impairs memory by attenuating mTORC1-dependent protein synthesis

Jennifer C. Tudor,¹ Emily J. Davis,^{1*} Lucia Peixoto,^{1†} Mathieu E. Wimmer,^{1‡} Erik van Tilborg,^{1§} Alan J. Park,^{1¶} Shane G. Poplawski,^{1||} Caroline W. Chung,¹ Robbert Havekes,^{1**} Jiayan Huang,² Evelina Gatti,^{3,4} Philippe Pierre,^{3,4} Ted Abel^{1††}

Sleep deprivation is a public health epidemic that causes wide-ranging deleterious consequences, including impaired memory and cognition. Protein synthesis in hippocampal neurons promotes memory and cognition. The kinase complex mammalian target of rapamycin complex 1 (mTORC1) stimulates protein synthesis by phosphorylating and inhibiting the eukaryotic translation initiation factor 4E-binding protein 2 (4EBP2). We investigated the involvement of the mTORC1-4EBP2 axis in the molecular mechanisms mediating the cognitive deficits caused by sleep deprivation in mice. Using an *in vivo* protein translation assay, we found that loss of sleep impaired protein synthesis in the hippocampus. Five hours of sleep loss attenuated both mTORC1-mediated phosphorylation of 4EBP2 and the interaction between eukaryotic initiation factor 4E (eIF4E) and eIF4G in the hippocampi of sleep-deprived mice. Increasing the abundance of 4EBP2 in hippocampal excitatory neurons before sleep deprivation increased the abundance of phosphorylated 4EBP2, restored the amount of eIF4E-eIF4G interaction and hippocampal protein synthesis to that seen in mice that were not sleep-deprived, and prevented the hippocampus-dependent memory deficits associated with sleep loss. These findings collectively demonstrate that 4EBP2-regulated protein synthesis is a critical mediator of the memory deficits caused by sleep deprivation.

INTRODUCTION

The consolidation of long-term hippocampus-dependent memory requires protein synthesis (1–3). The rate-limiting step for *de novo* protein production occurs at translation initiation, which is regulated by the insulin signaling pathway, including adenosine 5'-monophosphate-activated protein kinase (AMPK) and mammalian target of rapamycin (mTOR) (4–6). mTOR is a serine/threonine kinase that forms a complex called mTOR complex 1 (mTORC1) when associated with the regulatory-associated protein of mTOR (Raptor) (6–11). This complex releases the brake on protein synthesis initiation by mediating the phosphorylation of the translation repressor protein eukaryotic initiation factor 4E (eIF4E)-binding protein (4EBP). The predominant isoform of 4EBP in the brain

is 4EBP2 (12, 13). Phosphorylation of 4EBP2 prevents its binding to eIF4E, freeing eIF4E to stimulate translation. mTORC1 also promotes protein synthesis by activating the p70 S6 kinase (S6K) pathway (8–10).

Several studies have suggested that the mTOR signaling pathway is affected by sleep and sleep deprivation in various animal models. Sleep promotes the phosphorylation of 4EBP in the brains of cats (14), and periods of sleep have also been correlated with increased protein synthesis as measured by incorporation of radioactive leucine in the brains of rats (15) and nonhuman primates (16). In mice, we previously found that 5 hours of sleep deprivation leads to reduced abundance of mTOR in the hippocampus (17). Hippocampus-dependent memory is particularly sensitive to sleep loss (18–22). Here, we investigated whether sleep deprivation impairs protein synthesis *in vivo* and whether impaired protein synthesis causes the memory impairments associated with sleep deprivation.

RESULTS

Sleep deprivation impairs hippocampal protein synthesis *in vivo*

We first assessed whether hippocampal protein synthesis is reduced by sleep deprivation using a modified nonradioactive surface sensing of translation (SUnSET) assay that tags nascent proteins with puromycin, also known as ribopuromylation (23). Male C57BL/6J mice received one injection of puromycin (25 µg/µl) into the left lateral ventricle, and mice were then immediately sleep-deprived for 5 hours, whereas control mice were left undisturbed. Immediately after the sleep deprivation period, hippocampi were harvested, and the abundance of proteins tagged with puromycin was measured by Western blotting. Mice that had undergone 5 hours of sleep deprivation had a significantly reduced amount of puromycin-tagged proteins in the hippocampus compared to mice that had undisturbed sleep for 5 hours (Fig. 1A).

Our *in vivo* translation assay results were further corroborated when we compared the abundance of gene transcript to that of protein encoded

¹Department of Biology, School of Arts and Sciences, University of Pennsylvania, Philadelphia, PA 19104, USA. ²Global Statistical Science, Eli Lilly and Company, Lilly Corporate Center, Indianapolis, IN 46285, USA. ³Centre d'Immunologie de Marseille-Luminy, Aix Marseille Université UMR101, INSERM U1104, CNRS UMR7280, 13288 Marseille, France. ⁴Institute for Research in Biomedicine (iBIMED) and Aveiro Health Sciences Program, University of Aveiro, 3810-193 Aveiro, Portugal.

*Present address: Biomedical Sciences Graduate Program, University of California, San Francisco, San Francisco, CA 94158, USA.

†Present address: College of Medical Sciences, Washington State University, Spokane, WA 99202, USA.

‡Present address: Center for Neurobiology and Behavior, Department of Psychiatry, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA.

§Present address: Laboratory of Neuroimmunology and Developmental Origins of Disease, University Medical Center Utrecht, Lundlaan 6, 3584 GA Utrecht, Netherlands.

¶Present address: Department of Psychiatry, New York State Psychiatric Institute, Columbia University, New York, NY 10032, USA.

||Present address: Ibis Biosciences Inc. (an Abbott Company), Carlsbad, CA 92008, USA.

**Present address: Groningen Institute for Evolutionary Life Sciences, University of Groningen, 9747 AG Groningen, Netherlands.

††Corresponding author. Email: abele@sas.upenn.edu




Fig. 1. Sleep deprivation impairs hippocampal protein synthesis in vivo. (A) Western blotting analysis for puromycin, as a proxy for protein synthesis, in the hippocampus from non-sleep-deprived (NSD) or sleep-deprived (SD) male C57BL/6J mice injected intracerebroventricularly with puromycin. Puromycin signal was normalized to the glyceraldehyde-3-phosphate dehydrogenase (GAPDH) loading control and quantified (right panel). Data are means \pm SEM of 13 mice in each condition (* $P = 0.047$, t test). (B) Gene expression of *Arc* and *Hspa5/BiP* in hippocampal extracts from sleep-deprived mice relative to each in those from non-sleep-deprived mice, assessed by quantitative polymerase chain reaction (qPCR). Expression was normalized to that of housekeeping gene *Tuba4a*. Data are means \pm SEM of eight mice in each condition (** $P < 0.001$, t test). (C) Representative Western blots and quantitation of *Arc* and *Hspa5/BiP* abundance in hippocampal extracts. Abundance was normalized to the β -tubulin loading control. Data are means \pm SEM of 16 or 17 mice in each condition. All blots are representative of at least three independent assays. a.u., arbitrary units. The number of mice analyzed is noted in each bar.

by each of two genes in the hippocampus. The abundance of both *Arc* and *BiP* mRNA significantly increased about twofold after sleep deprivation (Fig. 1B), as previously shown (17), but we observed no commensurate increase in respective protein abundance (Fig. 1C). This dissociation between gene expression and protein abundance is consistent with a reduction of protein synthesis in the hippocampus after sleep deprivation.

Sleep deprivation reduces mTORC1 activity by increasing AMPK signaling in the hippocampus, which results in reduced 4EBP2 phosphorylation

Through bioinformatic analysis of microarray data, we previously showed that the group of genes for which expression is altered after sleep deprivation is enriched in genes that encode proteins in the insulin signaling pathway; these include AMPK and mTOR (17). mTOR and its multiple downstream effectors serve as a nexus for numerous cellular functions, including protein synthesis (8–10, 17). AMPK negatively regulates mTORC1 indirectly by phosphorylating and activating the tuberous sclerosis complex (TSC) (24) or directly by phosphorylating Raptor (Fig. 2) (25). To determine whether the insulin signaling pathway impairs protein synthesis after sleep deprivation, we deprived mice of sleep for 5 hours and probed the abundance of several insulin signaling pathway proteins in hippocampal extracts. We found that 5 hours of sleep deprivation significantly increased the phosphorylation of AMPK α in the hippocampus (Fig. 3A).




Fig. 2. Schematic of the signaling pathways regulating the initiation of protein synthesis. The insulin signaling pathway, which includes AMPK and mTOR, is one pathway that regulates protein synthesis initiation. When AMPK is activated, it inhibits mTORC1 activity through phosphorylation of either TSC or Raptor. The binding of mTOR and Raptor is necessary for mTORC1 formation and subsequent phosphorylation of p70 S6K or 4EBP2. Phosphorylation (activation) of S6K facilitates translation initiation by its phosphorylation (activation) of S6 ribosomal protein (S6). When 4EBP2 is phosphorylated (inhibited), eIF4E can form a complex with eIF4G to start cap-dependent translation.

mTOR forms at least two functionally distinct complexes that are mutually exclusive (26). mTORC1 is formed when mTOR is associated with Raptor, whereas mTORC2 is formed when rapamycin-insensitive companion of mTOR (Rictor) is associated with mTOR. We investigated whether 5 hours of sleep deprivation reduced mTORC1 or mTORC2 activity. To measure mTORC-specific activity, we examined the autophosphorylation of mTOR at Ser²⁴⁸¹ (27). We immunoprecipitated either Raptor or Rictor from hippocampal lysates and measured the amount of autophosphorylated mTOR that was bound to each. Blots from control pulldowns using rabbit immunoglobulin G (IgG) were negative for the proteins examined (fig. S2). We found that 5 hours of sleep deprivation specifically decreased the amount of autophosphorylated mTOR that was associated with Raptor (mTORC1) by 55% but not Rictor (mTORC2) (Fig. 3B). One possible mechanism by which increased AMPK activity would reduce mTOR activity is through phosphorylation of TSC2 (Fig. 2). We measured the phosphorylation of TSC2 at the AMPK phosphorylation site (Ser¹³⁸⁷) and did not find any changes in TSC2 phosphorylation in hippocampal extracts from sleep-deprived mice (fig. S3).

This reduction in mTORC1 but not mTORC2 activity led us to examine the downstream effectors of mTORC1-mediated translation, 4EBP2 and S6K (8). Sleep-deprived mice had significantly decreased amount of phosphorylated 4EBP2 in the hippocampus (Fig. 3C and fig. S4A). When mice were allowed to sleep for 2.5 hours after being sleep-deprived [a period of time sufficient for functional recovery from sleep deprivation (28, 29)], the amount of phosphorylated 4EBP2 was no longer significantly reduced (fig. S4B). In contrast, the abundance of phosphorylated S6K1 and phosphorylated S6 remained unchanged in the hippocampus from sleep-deprived mice (Fig. 3D). The phosphorylation of eIF2 α also did not change in sleep-deprived mice (fig. S5).




Fig. 3. Sleep deprivation affects AMPK-mTORC1-4EBP2 signaling pathway in the hippocampus. (A) Representative Western blots and quantitation of the ratios of phosphorylated to total AMPK α in hippocampal extracts from non-sleep-deprived mice and sleep-deprived mice. Data are means \pm SEM of six mice per condition ($*P = 0.032$, t test). (B) Representative Western blots and quantitation of hippocampal abundance of phosphorylated mTOR (p-mTOR) (Ser²⁴⁸¹) after immunoprecipitation (IP) with Raptor (left) or Rictor (right). Negative control blots for Rictor in the Raptor IP (left, below Raptor blot) and Raptor in the Rictor IP (right, below Rictor blot) are shown. Control input blots for phosphorylated mTOR (Ser²⁴⁸¹), Raptor, and Rictor are also shown. Data are means \pm SEM of 14 mice per condition ($*P = 0.042$, t test). (C and D) Representative Western blots and quantitation of phosphorylated 4EBP2 (p-4EBP2) (C) or phosphorylated S6K1 (p-S6K1) and phosphorylated S6 ribosomal protein (p-S6) (D) relative to their respective total protein abundance from hippocampus of non-sleep-deprived and sleep-deprived (after 5 hours) mice. β -Tubulin served as a loading control. Data are means \pm SEM of eight mice per condition ($***P < 0.005$, t test). (E) Representative Western blots and quantitation of hippocampal abundance of eIF4E after immunoprecipitation with eIF4G from non-sleep-deprived and sleep-deprived mice. Control input blots for eIF4E and eIF4G are also shown. Data are means \pm SEM of eight mice per condition ($*P = 0.047$, t test). All blots are representative of at least three independent assays. The number of mice analyzed is noted in each bar.

Reduced mTORC1 activity and increased 4EBP2 phosphorylation observed in the hippocampus of sleep-deprived mice would suggest that eIF4E binding to eIF4G would be attenuated (Fig. 2). We immunoprecipitated eIF4G and measured the amount of bound eIF4E. We found that 5 hours of sleep deprivation in mice decreased the eIF4E-eIF4G association in the hippocampus (Fig. 3E). Thus, sleep deprivation attenuates protein synthesis initiation by specifically affecting the AMPK-mTORC1-4EBP2 signaling pathway, resulting in decreased eIF4E-eIF4G association in the hippocampus.

Viral expression of 4EBP2 in the hippocampus prevents deficits in protein synthesis and memory impairment caused by sleep deprivation

Given the decrease in 4EBP2 phosphorylation after 5 hours of sleep deprivation, we examined whether the viral expression of 4EBP2 might rescue memory deficits caused by sleep deprivation. We expressed 4EBP2 in hippocampal excitatory neurons of adult male C57BL/6J mice using a calcium/calmodulin-dependent protein kinase II α (CaMKII α) promoter fragment in adeno-associated viruses (AAVs) (Fig. 4A). Three weeks after AAV vector injection into the hippocampus, phosphorylated 4EBP2 was significantly increased fourfold compared to total 4EBP2 in the hippocampus of mice injected with the 4EBP2 AAV (Fig. 4B). We also confirmed that viral expression was limited to excitatory neurons by assessing colabeling of CaMKII α and hemagglutinin (HA)-tagged 4EBP2 (4EBP2-HA) in hippocampal cells in situ (Fig. 4C). Viral expression of 4EBP2-HA did not colabel with parvalbumin (Fig. 4D) or glial fibrillary acidic protein (GFAP; Fig. 4E), demonstrating that 4EBP2-HA was not expressed in parvalbumin-positive inhibitory neurons or glial cells.

To determine whether viral expression of 4EBP2 rescues the reduced interaction between eIF4E and eIF4G after sleep deprivation, we immunoprecipitated eIF4G and examined eIF4E abundance bound to eIF4G in undisturbed and sleep-deprived mice injected with control enhanced green fluorescent protein (eGFP) AAV or 4EBP2 AAV. We found that expression of 4EBP2 in the hippocampus rescued the abundance of eIF4E bound to eIF4G after sleep deprivation (Fig. 4F). Using the *in vivo* translation assay described above, we measured the amount of puromycin incorporation into proteins in the hippocampus after 5 hours of sleep deprivation in mice virally expressing 4EBP2 or eGFP (control) for 3 weeks. AAV-induced increase of 4EBP2 expression in the hippocampus prevented the reduced protein synthesis that was caused by sleep deprivation (Fig. 4G). Three weeks of 4EBP2 viral expression were sufficient to restore protein synthesis amounts in the hippocampus of sleep-deprived mice. Therefore, we performed subsequent behavioral experiments at this time point after virus injection.

We previously showed that 5 hours of sleep deprivation causes hippocampus-dependent memory deficits (21, 28, 30). Here, we investigated whether restoring protein synthesis is sufficient to prevent the memory deficits associated with sleep deprivation.


We trained mice expressing 4EBP2 AAV or eGFP AAV in the hippocampus in an object-place recognition task (Fig. 4H). This spatial task exploits the preference for novelty in mice (31, 32). During training, mice explored and learned the location of three objects within an arena. Immediately after training, mice expressing 4EBP2 virus or control eGFP virus were either sleep-deprived for 5 hours or left undisturbed. The next day, mice were tested for object-location memory. Consistent with previous studies, mice expressing eGFP that were left undisturbed preferentially explored the displaced object, which demonstrates consolidated memory (30).

Fig. 4. Viral expression of 4EBP2 in the hippocampus prevents deficits in protein synthesis and memory impairment caused by sleep deprivation. (A) Schematics of the pAAV₉-CaMKII α 0.4-eGFP or pAAV₉-CaMKII α 0.4-eIF4EBP2-HA vectors used to express eGFP or 4EBP2 in hippocampal excitatory neurons in mice. ITR, inverted terminal repeat. (B) Representative Western blots and quantitation of the ratio between phosphorylated 4EBP2 and total 4EBP2 in the hippocampus of mice injected with either the eGFP or the 4EBP2 vector. Abundance was normalized to GAPDH loading control. Data are means \pm SD from six mice in each condition (** $P < 0.005$, t test). (C to E) Immunofluorescence for 4EBP2-HA (green), CaMKII (red), parvalbumin (Pvalb; red), or GFAP (red) in the CA3 region of the hippocampus from 4EBP2-HA-injected mice. Arrows indicate colocalization. Scale bar, 50 μ m. Images represent four mice per condition (three to five images per mouse). (F) Representative Western blots and quantitation of hippocampal abundance of eIF4E after immunoprecipitation with eIF4G from eGFP- and 4EBP2-HA-injected mice that were either non-sleep-deprived or sleep-deprived. Control input blots for eIF4E and eIF4G are also shown. Data are means \pm SEM of six mice per condition [sleep deprivation effect: $F_{1,20} = 9.856$, ** $P = 0.0052$; virus effect: $F_{1,20} = 16.53$, *** $P = 0.0006$; interaction effect: $F_{1,20} = 6.988$, * $P = 0.0156$; two-way analysis of variance (ANOVA)]. (G) Representative Western blots and quantitation of proteins labeled with puromycin from hippocampi of non-sleep-deprived and sleep-deprived mice 3 weeks after being injected with eGFP or 4EBP2. Abundance was normalized to GAPDH loading control. Data are means \pm SEM of six or eight mice in each condition [sleep deprivation effect: $F_{1,22} = 1.585$, $P = 0.221$; virus effect: $F_{1,22} = 0.577$, $P = 0.456$; interaction effect: $F_{1,22} = 5.201$, * $P = 0.033$; two-way ANOVA]. (H) Performance of mice expressing eGFP or 4EBP2 in a hippocampus-dependent object-place recognition task (schematic, top) when either sleep-deprived for 5 hours immediately after training or left undisturbed. Data are means \pm SEM of six or seven mice in each condition. (eGFP, * $P < 0.013$; 4EBP2, $P = 0.149$; Wilcoxon rank-sum test). Dotted line indicates chance performance. All blots are representative of at least three independent assays. The number of mice analyzed is noted in each bar.

Sleep-deprived mice expressing eGFP did not exhibit increased exploration of the displaced object, demonstrating a failure to consolidate memory of it. Mice expressing 4EBP2 AAV preferentially explored the displaced object, regardless of whether they were sleep-deprived or not (Fig. 4H), indicating that 4EBP2 AAV mice were resistant to memory deficits caused by sleep deprivation. Collectively, these data demonstrate that increased 4EBP2 expression in hippocampal excitatory neurons significantly increased phosphorylated 4EBP2 abundance, which restored protein synthesis in the hippocampus and prevented memory deficits caused by sleep deprivation.

DISCUSSION

Previous work suggested that sleep deprivation affects protein synthesis signaling pathways, specifically the insulin signaling pathway (17). Here, we found that hippocampus-dependent memory impairments caused by sleep deprivation are due to impaired protein synthesis initiation in the hippocampus (summarized in fig. S1). A specific subset of the insulin signaling pathway—AMPK-mTOR-4EBP2 signaling—was affected by sleep deprivation, which was sufficient to cause hippocampus-dependent memory deficits in mice. Furthermore, we rescued these memory def-



icits when we restored protein synthesis in the hippocampus of these mice with 4EBP2 AAV infection, which increased 4EBP2 phosphorylation in hippocampal excitatory neurons. These findings suggest that sleep deprivation impairs memory by reducing protein synthesis.

Our study found that sleep deprivation impairs protein synthesis necessary for memory consolidation. Our work supports the hypothesis that one of the functions of sleep is to facilitate protein synthesis (33, 34). We demonstrated that sleep deprivation impaired protein synthesis in the hippocampus using a nonradioactive *in vivo* translation assay, which was predicted by our previous findings that 5 hours of sleep deprivation lowers the abundance of mTOR in the hippocampus (17). This is consistent with previous findings that rapamycin administration inhibiting mTORC1 impairs sleep-dependent cortical plasticity (14). Grønli and colleagues (35) have also shown that translation factors are affected by sleep deprivation. Conversely, periods of sleep have been correlated with increased protein synthesis as measured by a radioactive translation assay in other organisms (15, 16). Furthermore, sleep promotes the phosphorylation of 4EBP2 (14), which is in line with our findings that phosphorylated 4EBP2 is reduced with sleep deprivation. Our study demonstrates the functional importance of the impact of sleep deprivation on protein synthesis because we show that rescuing protein synthesis is sufficient to prevent these sleep deprivation-induced memory impairments. Our rescue experiments

demonstrate that it is the impact of sleep deprivation on protein synthesis in the hippocampus that is critical for spatial memory impairments. In future experiments, it would be interesting to see whether short periods of sleep deprivation affect other behaviors by attenuating protein synthesis in other brain regions.

We found that sleep deprivation specifically attenuated mTORC1 activity, which reduced the abundance of phosphorylated 4EBP2, but not of phosphorylated S6K, in the mouse hippocampus. This result, although surprising, is not unique to sleep deprivation. Neural stem cell self-renewal is also regulated by the specific targeting of mTORC1 to 4EBP2, and not S6K (36). One possible explanation for the specificity of the impact of sleep deprivation on the 4EBP2 pathway but not S6K signaling may be due to the action of specific protein phosphatases. Our previously published microarray data revealed several phosphatase genes with increased expression in the hippocampus after sleep deprivation (17). It would be interesting to examine in the future the role of phosphatases after sleep deprivation to discern the mechanism by which sleep deprivation selectively affects the phosphorylation of 4EBP2. Another possible hypothesis that explains why sleep deprivation specifically affects 4EBP2 activity may be due to its association with Raptor.

Previous work in *Aplysia* has shown that Raptor signaling and 4EBP2 signaling are tightly regulated (37). We restored the amount of protein synthesis in sleep-deprived mice to that seen in non-sleep-deprived mice using 4EBP2 AAV infection into the hippocampus. Thus, increasing the abundance of 4EBP2 may increase the association between Raptor and 4EBP2. It is important to note that 4EBP2 undergoes an alternate post-translational modification other than phosphorylation. 4EBP2 deamidation occurs when two C-terminal asparagines spontaneously convert to aspartates (38). Sleep deprivation may increase 4EBP2 deamidation in the hippocampus, and increasing 4EBP2 expression would increase the Raptor-4EBP2 association. This increase in Raptor-4EBP2 interaction, with reduced 4EBP2 sequestration of eIF4E, would ultimately decrease translation repression (39). Our finding that 4EBP2 AAV infection rescued the abundance of eIF4E bound to eIF4G supports this hypothesis. These results suggest that sleep deprivation may be affecting Raptor because of its known interaction with 4EBP2. Additionally, we did not find changes to TSC2 phosphorylation as a result of sleep deprivation. This finding suggests that AMPK signaling inhibits mTORC1 activity through Raptor phosphorylation. Future experiments could probe the involvement of Raptor in the insulin signaling pathway and its changes after sleep deprivation.

We did not see increased protein synthesis or enhanced memory in mice expressing the 4EBP2 virus that were not sleep-deprived. We speculate that we observed a ceiling effect, where depressed translation could be rescued by 4EBP2 expression, but translation could not be facilitated beyond physiological abundance by 4EBP2 expression. The specificity of restoring memory in sleep-deprived mice clearly demonstrated the importance of AMPK-mTORC1-4EBP2 activity for protein synthesis and subsequent memory impairment after sleep deprivation.

We showed that rescuing 4EBP2 expression and subsequent protein synthesis was sufficient to prevent the memory deficits associated with sleep deprivation. We have previously shown that cyclic AMP (cAMP) signaling is also impaired with sleep deprivation, leading to cognitive deficits (28, 30). Although it is possible that, with sleep deprivation, cAMP signaling and mTOR signaling occur in parallel and independent of one another, there are points of interaction between the cAMP and mTOR signaling pathways. One way in which sleep deprivation affects both cAMP and mTOR signaling may be through phosphodiesterase 4D (PDE4D), a protein that degrades cAMP. PDE4D can bind to RAS homolog enriched in brain (Rheb), inhibiting Rheb from activating mTORC1 (40). Thus, one possible explanation is that sleep deprivation reduces cAMP,

which would elicit more PDE4D interaction with Rheb and thereby decrease the interaction between Rheb and mTOR (40). This would ultimately lead to reduced mTORC1-mediated translation. Previously, we have shown that sleep deprivation specifically increases PDE4A5, not PDE4D (28, 30). It would be very interesting to examine in future experiments whether PDE4A5 interacts with Rheb in a similar fashion as PDE4D, as this has yet to be determined (41).

The spatial, contextual, and declarative memories that are dependent on the hippocampus are particularly susceptible to the effects of insufficient sleep (18, 19, 42). The object-based location memory task we used to measure the impact of sleep deprivation on spatial memory in mice is specifically dependent on the hippocampus. Therefore, we focused our studies on this region of the brain. Others have demonstrated that sleep deprivation also increases AMPK activity in the hypothalamus and the basal forebrain (43, 44). Further, they have shown that this increase in AMPK activity is accompanied by decreased adenosine 5'-triphosphate (ATP) abundance (44). Protein synthesis is an energy-rich process, and the insulin signaling pathway is central to regulating whether protein synthesis occurs given the metabolic resources available to the cell (24). Increased AMPK activity with sleep deprivation suggests ATP depletion in the hippocampus (38), which would lead to reductions in protein synthesis and impairments in memory consolidation. Many of the other molecular and cellular consequences of sleep deprivation in the hippocampus including decreased cAMP signaling (22, 28, 30) and increased adenosine (22, 45, 46) could be due to reduced abundance of ATP. Thus, our results support the hypothesis that one of the functions of sleep is cellular energy restoration (47).

MATERIALS AND METHODS

Animal subjects

Male C57BL/6J mice were obtained at 6 to 8 weeks of age from The Jackson Laboratory for these experiments. Mice were housed individually on a 12-hour light/dark schedule with lights on at 7:00 a.m. Food and water were available ad libitum. All experiments were conducted according to the National Institutes of Health *Guide for the Care and Use of Laboratory Animals* and were approved by the University of Pennsylvania's Institutional Animal Care and Use Committee.

Sleep deprivation

Mice were sleep-deprived for 5 hours using the gentle handling method (17, 28, 31, 48–50). Briefly, this technique is composed of manual cage tapping, cage jostling, and nestlet disturbance. Before sleep deprivation, each animal was handled daily for 3 to 6 days using the same interventions used during sleep deprivation for 1 to 2 min. Non-sleep-deprived mice were left undisturbed in their home cages.

Western blotting and enzyme-linked immunosorbent assay

Mice were sacrificed by cervical dislocation at the conclusion of sleep deprivation experiments. The hippocampus was rapidly dissected and flash-frozen on dry ice. Total protein lysates were prepared by homogenizing the hippocampus in buffer containing 50 mM tris-HCl (pH 7.5), 150 mM KCl, 1 mM EDTA, 1 mM dithiothreitol, as well as protease (Roche) and phosphatase inhibitors (Thermo Scientific). For Western blotting, samples were normalized on the basis of total protein content measured by Bradford assay (Bio-Rad). Proteins were separated by 4 to 20% tris-glycine SDS-polyacrylamide gel electrophoresis (Bio-Rad) and transferred to polyvinylidene difluoride membranes. Membranes were blocked in 5% bovine

serum albumin–tris-buffered saline with Tween 20 (BSA-TBST) or 5% milk + 0.5% BSA-TBST and incubated overnight at 4°C with primary antibody. Antibody concentrations were at the manufacturers' recommendations [p-4EBP, 4EBP2, AMPK α , Bip, eIF4E, eIF4G1, p-mTOR, mTOR, p-S6, S6, p-S6K, Rictor, p-TSC2, and TSC2 (1:1000; Cell Signaling); p-AMPK α and S6K (1:2000; Cell Signaling); p-eIF2 α (1:1000; Invitrogen); Raptor (1:2000; Bethyl Laboratories); Arc (1:2000; Synaptic Systems); and puromycin 12D10 (1:5000; Millipore) from stock (5 mg/ml)].

Afterward, they were washed and incubated with appropriate horseradish peroxidase–conjugated goat anti-mouse or anti-rabbit IgG (1:5000; Santa Cruz Biotechnology) at room temperature for 1 hour. Blots were washed and exposed via chemiluminescence using ImageQuant LAS 4000 (GE) and quantified using ImageJ. The density of signal was normalized to GAPDH (1:1000; Santa Cruz Biotechnology) or β -tubulin (1:10,000; Sigma-Aldrich). For immunoprecipitations, hippocampal lysates were precleared using protein A/G agarose beads (Thermo Scientific/Pierce) while rotating for 1 hour at 4°C. Antibody was added to the cell lysates and incubated while rotating overnight at 4°C. Protein A/G agarose beads were then added to the lysates and incubated while rotating overnight at 4°C. Lysates were centrifuged and agarose beads were washed four times in lysis buffer. Immunoprecipitates were analyzed by Western blot. Negative control IgG pulldowns were analyzed in lysates that were treated with rabbit IgG (Bethyl Laboratories). Enzyme-linked immunosorbent assay (ELISA) analysis of hippocampal lysates was conducted in accordance to the manufacturer's guidelines (Cell Signaling).

Quantitative PCR

RNA preparation, complementary DNA (cDNA) synthesis, and qPCR analyses were performed as previously described (17, 51). Briefly, RNA concentration and purity were quantified by NanoDrop spectrophotometer (Thermo Scientific), and cDNA was generated using the RETROscript Kit (Ambion) with 1 μ g of RNA as template. Three technical replicates were measured by qPCR using ABI Prism 7000. Data were normalized to *Actg1*, *Hprt*, and *Tuba4a*. Fold change was calculated from the ΔC_t values with corrections for housekeeping gene expression for each sample.

In vivo translation assay

Proteins were labeled using an adaptation of the SUNSET protocol (52–54), also known as ribopuromycylation (23). A week before sleep deprivation experiments to allow for postoperative recovery, mice were anesthetized with isoflurane and mounted onto a stereotaxic apparatus. Intracerebroventricular cannulae were implanted unilaterally at –0.5-mm anteroposterior and +1.0-mm mediolateral from bregma. At the start of sleep deprivation experiments, 25 μ g in 1 μ l of puromycin (Sigma-Aldrich) was infused into the ventricle over 5 min. At the end of sleep deprivation experiments, the contralateral hippocampus was dissected and flash-frozen. Proteins labeled with puromycin were identified by Western blot analysis from hippocampal lysates.

Viral constructs and surgeries

The eIF4EBP2 gene construct was generated using GeneArt (Millipore) with an HA peptide tag at the C-terminal end. A 0.4-kb CaMKII α promoter fragment was used to restrict expression to excitatory neurons. The pAAV $_9$ -CaMKII α 0.4-eIF4EBP2-HA recombinant, along with control pAAV $_9$ -CaMKII α 0.4-eGFP, was constructed and packaged by the University of Pennsylvania Vector Core. Viral titers ranged from 2.4×10^{12} to 1.47×10^{13} genome copy per microliter. Mice were anesthetized with isoflurane and mounted onto a stereotaxic apparatus. Depending on the titer, about

1 μ l of the virus suspension was injected bilaterally at –1.9-mm anteroposterior, \pm 1.5-mm mediolateral, and –1.5-mm dorsoventral from bregma into the hippocampi of mice using a 33-gauge beveled NanoFil needle, a microsyringe pump, and controller (WPI). Three weeks after surgery, mice were tested in the object-place recognition task and hippocampal *in vivo* translation was assayed.

Perfusion and immunohistochemistry

Mice were transcardially perfused with 10 ml of phosphate-buffered saline (PBS) before infusion of 4% paraformaldehyde (PFA) in PBS. Perfused brains were postfixed for 24 hours in 4% PFA at 4°C before immersion in 30% sucrose. After cryoprotection with 30% sucrose, 30- μ m coronal sections were cut. To validate expression of 4EBP2 virus in the hippocampus, immunohistochemical analyses were performed on mouse brain sections. Sections were washed three times in PBS for 5 min each and then incubated with 0.3% H $_2$ O $_2$ in PBS for 30 min. Afterward, sections were washed three times in PBS for 10 min each. They were blocked in 5% of appropriate serum (normal goat serum and normal donkey serum) in 0.1% Triton X-100 in PBS for 30 min. Sections were incubated with primary antibodies [anti-HA rat monoclonal antibody (1:500; Roche), anti-CaMKII goat polyclonal (1:50; Santa Cruz Biotechnology), anti-parvalbumin rabbit polyclonal (1:1000; Abcam), and anti-GFAP mouse polyclonal (1:500; EMD Millipore)] in 1% serum in 1% Triton X-100/PBS at room temperature overnight. Sections were washed three times in PBS for 5 min each. They were incubated with appropriate secondary antibodies (1:1000) (goat anti-rat IgG, Alexa Fluor 488; donkey anti-goat IgG, Alexa Fluor 555; donkey anti-rabbit IgG, Alexa Fluor 555; and goat anti-mouse IgG, Alexa Fluor 555) in PBS for 4 hours at room temperature in the dark. They were washed six times over 60 min in PBS and mounted onto slides using 0.7% gelatin in H $_2$ O. Slides were allowed to dry overnight and then coverslipped with PermaFluor. Sections were imaged with an epifluorescent microscope to determine specificity of viral 4EBP2 expression in the hippocampus.

Object-place recognition task

This hippocampus-dependent memory task was conducted as previously described (30, 31, 55, 56). Sixteen days after viral surgeries, mice were handled for 2 min each day for five consecutive days leading up to experimentation. At the beginning of the light phase (zeitgeber time 0), mice were placed in the empty box for 6 min for habituation. After a 3-min intertrial interval, mice were placed in the box with three different objects (a 100-ml glass bottle, a white cylinder, and a metallic rectangular tower) for three consecutive 6-min training sessions. Each training session was separated by a 3-min interval during which the animals were returned to the holding cages. Directly after training, mice were either sleep-deprived or left undisturbed. Twenty-four hours after the training session, mice were reintroduced to the spatial context in a single 6-min test session with one of three objects moved to a novel location. Objects and locations were balanced between treatment groups. A video of behavior was recorded and scored offline by an experimenter that was blind to treatment. Exploration of the objects was defined as the amount of time mice were oriented toward an object within close proximity or touching it.

Statistical analyses

All blots and images are representative of at least three independent assays. Data represent means \pm SEM. Student's *t* tests were used to analyze the gene expression and biochemical data from sleep deprivation experiments. Two-way ANOVAs were used to analyze the biochemical data from the four-group viral experiments. Dunnett's test was used for post hoc comparisons. The Wilcoxon rank-sum test and the Kruskal-Wallis test were used to

analyze the behavioral data. When $P < 0.05$, differences were considered statistically significant.

SUPPLEMENTARY MATERIALS

www.sciencesignaling.org/cgi/content/full/9/425/ra41/DC1

Fig. S1. Schematic showing the mechanism of attenuated translation in the hippocampus caused by sleep deprivation.

Fig. S2. Control immunoblots of hippocampal extracts treated with rabbit IgG.

Fig. S3. The abundance of phosphorylated TSC2 does not change after 5 hours of sleep deprivation.

Fig. S4. Five hours of sleep deprivation reduces the abundance of phosphorylated 4EBP2, which rebounds after 2.5 hours of recovery sleep.

Fig. S5. Abundance of phosphorylated eIF2 α does not change after 5 hours of sleep deprivation.

REFERENCES AND NOTES

1. R. Bourchouladze, T. Abel, N. Berman, R. Gordon, K. Lapidus, E. R. Kandel, Different training procedures recruit either one or two critical periods for contextual memory consolidation, each of which requires protein synthesis and PKA. *Learn. Mem.* **5**, 365–374 (1998).
2. W. C. Abraham, J. M. Williams, LTP maintenance and its protein synthesis-dependence. *Neurobiol. Learn. Mem.* **89**, 260–268 (2008).
3. R. J. Kelleher III, A. Govindarajan, S. Tonegawa, Translational regulatory mechanisms in persistent forms of synaptic plasticity. *Neuron* **44**, 59–73 (2004).
4. T. E. Dever, Gene-specific regulation by general translation factors. *Cell* **108**, 545–556 (2002).
5. F. Gebauer, M. W. Hentze, Molecular mechanisms of translational control. *Nat. Rev. Mol. Cell Biol.* **5**, 827–835 (2004).
6. S. A. Buffington, W. Huang, M. Costa-Mattoli, Translational control in synaptic plasticity and cognitive dysfunction. *Annu. Rev. Neurosci.* **37**, 17–38 (2014).
7. J. O. Lipton, M. Sahin, The neurology of mTOR. *Neuron* **84**, 275–291 (2014).
8. N. Hay, N. Sonenberg, Upstream and downstream of mTOR. *Genes Dev.* **18**, 1926–1945 (2004).
9. Q. Yang, K.-L. Guan, Expanding mTOR signaling. *Cell Res.* **17**, 666–681 (2007).
10. C. A. Hoeffer, E. Klann, mTOR signaling: At the crossroads of plasticity, memory and disease. *Trends Neurosci.* **33**, 67–75 (2010).
11. V. Albert, M. N. Hall, mTOR signaling in cellular and organismal energetics. *Curr. Opin. Cell Biol.* **33**, 55–66 (2015).
12. K. Tsukiyama-Kohara, F. Poulin, M. Kohara, C. T. DeMaria, A. Cheng, Z. Wu, A.-C. Gingras, A. Katsume, M. Elchebly, B. M. Spiegelman, M.-E. Harper, M. L. Tremblay, N. Sonenberg, Adipose tissue reduction in mice lacking the translational inhibitor 4E-BP1. *Nat. Med.* **7**, 1128–1132 (2001).
13. J. L. Banko, F. Poulin, L. Hou, C. T. DeMaria, N. Sonenberg, E. Klann, The translation repressor 4E-BP2 is critical for eIF4F complex formation, synaptic plasticity, and memory in the hippocampus. *J. Neurosci.* **25**, 9581–9590 (2005).
14. J. Seibt, M. C. Dumoulin, S. J. Aton, T. Coleman, A. Watson, N. Naidoo, M. G. Frank, Protein synthesis during sleep consolidates cortical plasticity in vivo. *Curr. Biol.* **22**, 676–682 (2012).
15. P. Ramm, C. T. Smith, Rates of cerebral protein synthesis are linked to slow wave sleep in the rat. *Physiol. Behav.* **48**, 749–753 (1990).
16. H. Nakanishi, Y. Sun, R. K. Nakamura, K. Mori, M. Ito, S. Suda, H. Namba, F. I. Storch, T. P. Dang, W. Mendelson, M. Mishkin, C. Kennedy, J. C. Gillin, C. B. Smith, L. Sokoloff, Positive correlations between cerebral protein synthesis rates and deep sleep in *Macaca mulatta*. *Eur. J. Neurosci.* **9**, 271–279 (1997).
17. C. G. Vecsey, L. Peixoto, J. H. Choi, M. Wimmer, D. Jaganath, P. J. Hernandez, J. Blackwell, K. Meda, A. J. Park, S. Hannehalli, T. Abel, Genomic analysis of sleep deprivation reveals translational regulation in the hippocampus. *Physiol. Genomics* **44**, 981–991 (2012).
18. T. Abel, R. Havekes, J. M. Saletin, M. P. Walker, Sleep, plasticity and memory from molecules to whole-brain networks. *Curr. Biol.* **23**, R774–R788 (2013).
19. S. Banks, D. F. Dinges, Behavioral and physiological consequences of sleep restriction. *J. Clin. Sleep Medicine* **3**, 519–528 (2007).
20. G. Curcio, M. Ferrara, L. De Gennaro, Sleep loss, learning capacity and academic performance. *Sleep Med. Rev.* **10**, 323–337 (2006).
21. L. A. Graves, E. A. Heller, A. I. Pack, T. Abel, Sleep deprivation selectively impairs memory consolidation for contextual fear conditioning. *Learn. Mem.* **10**, 168–176 (2003).
22. R. Havekes, C. G. Vecsey, T. Abel, The impact of sleep deprivation on neuronal and glial signaling pathways important for memory and synaptic plasticity. *Cell. Signal.* **24**, 1251–1260 (2012).
23. A. David, B. P. Dolan, H. D. Hickman, J. J. Knowlton, G. Clavarino, P. Pierre, J. R. Bennink, J. W. Yewdell, Nuclear translation visualized by ribosome-bound nascent chain puromycylation. *J. Cell Biol.* **197**, 45–57 (2012).
24. K. Inoki, J. Kim, K.-L. Guan, AMPK and mTOR in cellular energy homeostasis and drug targets. *Annu. Rev. Pharmacol. Toxicol.* **52**, 381–400 (2012).
25. D. M. Gwinn, D. B. Schackelford, D. F. Egan, M. M. Mihaylova, A. Mery, D. S. Vasquez, B. E. Turk, R. J. Shaw, AMPK phosphorylation of raptor mediates a metabolic checkpoint. *Mol. Cell* **30**, 214–226 (2008).
26. D. D. Sarbassov, S. M. Ali, D.-H. Kim, D. A. Guertin, R. R. Latek, H. Erdjument-Bromage, P. Tempst, D. M. Sabatini, Rictor, a novel binding partner of mTOR, defines a rapamycin-insensitive and raptor-independent pathway that regulates the cytoskeleton. *Curr. Biol.* **14**, 1296–1302 (2004).
27. G. A. Soliman, H. A. Acosta-Jaquez, E. A. Dunlop, B. Ekim, N. E. Maj, A. R. Tee, D. C. Fingar, mTOR Ser-2481 autophosphorylation monitors mTORC-specific catalytic activity and clarifies rapamycin mechanism of action. *J. Biol. Chem.* **285**, 7866–7879 (2010).
28. C. G. Vecsey, G. S. Baillie, D. Jaganath, R. Havekes, A. Daniels, M. Wimmer, T. Huang, K. M. Brown, X.-Y. Li, G. Descalzi, S. S. Kim, T. Chen, Y.-Z. Shang, M. Zhuo, M. D. Houslay, T. Abel, Sleep deprivation impairs cAMP signalling in the hippocampus. *Nature* **461**, 1122–1125 (2009).
29. R. Huber, T. DeBoer, I. Tobler, Effects of sleep deprivation on sleep and sleep EEG in three mouse strains: Empirical data and simulations. *Brain Res.* **857**, 8–19 (2000).
30. R. Havekes, V. M. Bruinenberg, J. C. Tudor, S. L. Ferri, A. Baumann, P. Meerlo, T. Abel, Transiently increasing cAMP levels selectively in hippocampal excitatory neurons during sleep deprivation prevents memory deficits caused by sleep loss. *J. Neurosci.* **34**, 15715–15721 (2014).
31. T.-M. Prince, M. Wimmer, J. Choi, R. Havekes, S. Aton, T. Abel, Sleep deprivation during a specific 3-hour time window post-training impairs hippocampal synaptic plasticity and memory. *Neurobiol. Learn. Mem.* **109**, 122–130 (2014).
32. S. G. Poplawski, H. Schoch, M. E. Wimmer, J. D. Hawk, J. L. Walsh, K. P. Giese, T. Abel, Object-location training elicits an overlapping but temporally distinct transcriptional profile from contextual fear conditioning. *Neurobiol. Learn. Mem.* **116**, 90–95 (2014).
33. J. P. Wisor, The sleep-deprived hippocampus: A loss in translation. *Physiol. Genomics* **45**, 26–27 (2013).
34. J. Grønli, J. Soulé, C. R. Bramham, Sleep and protein synthesis-dependent synaptic plasticity: Impacts of sleep loss and stress. *Front. Behav. Neurosci.* **7**, 224 (2014).
35. J. Grønli, G. Dagestad, A. M. Milde, R. Murison, C. R. Bramham, Post-transcriptional effects and interactions between chronic mild stress and acute sleep deprivation: Regulation of translation factor and cytoplasmic polyadenylation element-binding protein phosphorylation. *Behav. Brain Res.* **235**, 251–262 (2012).
36. N. W. Hartman, T. V. Lin, L. Zhang, G. E. Paquelet, D. M. Feliciano, A. Bordey, mTORC1 targets the translational repressor 4E-BP2, but not S6 kinase 1/2, to regulate neural stem cell self-renewal in vivo. *Cell Rep.* **5**, 433–444 (2013).
37. M. Carroll, J. Dyer, W. S. Sossin, Serotonin increases phosphorylation of synaptic 4EBP through TOR, but eukaryotic initiation factor 4E levels do not limit somatic cap-dependent translation in *aplysia* neurons. *Mol. Cell Biol.* **26**, 8586–8598 (2006).
38. M. Bidnost, I. Ran, M. R. Sanchez-Carbente, Y. Martineau, A.-C. Gingras, C. Gkogkas, B. Rought, C. R. Bramham, W. S. Sossin, M. Costa-Mattoli, L. DesGroseillers, J.-C. Lacaille, N. Sonenberg, Postnatal deamidation of 4E-BP2 in brain enhances its association with raptor and alters kinetics of excitatory synaptic transmission. *Mol. Cell* **37**, 797–808 (2010).
39. T. E. Graber, P. K. McCarroll, W. S. Sossin, A recollection of mTOR signaling in learning and memory. *Learn. Mem.* **20**, 518–530 (2013).
40. H. W. Kim, S. H. Ha, M. N. Lee, E. Huston, D.-H. Kim, S. K. Jang, P.-G. Suh, M. D. Houslay, S. H. Ryu, Cyclic AMP controls mTOR through regulation of the dynamic interaction between Rheb and phosphodiesterase 4D. *Mol. Cell Biol.* **30**, 5406–5420 (2010).
41. D. H. Maurice, H. Ke, F. Ahmad, Y. Wang, J. Chung, V. C. Manganiello, Advances in targeting cyclic nucleotide phosphodiesterases. *Nat. Rev. Drug Discov.* **13**, 290–314 (2014).
42. S. Diekelmann, J. Born, The memory function of sleep. *Nat. Rev. Neurosci.* **11**, 114–126 (2010).
43. S. Chikahisa, N. Fujiki, K. Kitaoka, N. Shimizu, H. Séi, Central AMPK contributes to sleep homeostasis in mice. *Neuropharmacology* **57**, 369–374 (2009).
44. M. Dworak, R. W. McCarley, T. Kim, A. V. Kalinchuk, R. Basheer, Sleep and brain energy levels: ATP changes during sleep. *J. Neurosci.* **30**, 9007–9016 (2010).
45. C. Florian, C. G. Vecsey, M. M. Halassa, P. G. Haydon, T. Abel, Astrocyte-derived adenosine and A₁ receptor activity contribute to sleep loss-induced deficits in hippocampal synaptic plasticity and memory in mice. *J. Neurosci.* **31**, 6956–6962 (2011).
46. M. M. Halassa, C. Florian, T. Fellin, J. R. Muñoz, S.-Y. Lee, T. Abel, P. G. Haydon, M. G. Frank, Astrocytic modulation of sleep homeostasis and cognitive consequences of sleep loss. *Neuron* **61**, 213–219 (2009).
47. S. Chikahisa, H. Séi, The role of ATP in sleep regulation. *Front. Neurol.* **2**, 87 (2011).
48. R. Hagewoud, R. Havekes, A. Novati, J. N. Keijsers, E. A. Van der Zee, P. Meerlo, Sleep deprivation impairs spatial working memory and reduces hippocampal AMPA receptor phosphorylation. *J. Sleep Res.* **19**, 280–288 (2010).
49. R. Hagewoud, S. N. Whitcomb, A. N. Heeringa, R. Havekes, J. M. Koolhaas, P. Meerlo, A time for learning and a time for sleep: The effect of sleep deprivation on contextual fear conditioning at different times of the day. *Sleep* **33**, 1315–1322 (2010).

50. C. G. Vecsey, M. E. J. Wimmer, R. Havekes, A. J. Park, I. J. Perron, P. Meerlo, T. Abel, Daily acclimation handling does not affect hippocampal long-term potentiation or cause chronic sleep deprivation in mice. *Sleep* **36**, 601–607 (2013).
51. C. G. Vecsey, J. D. Hawk, K. M. Lattal, J. M. Stein, S. A. Fabian, M. A. Attner, S. M. Cabrera, C. B. McDonough, P. K. Brindle, T. Abel, M. A. Wood, Histone deacetylase inhibitors enhance memory and synaptic plasticity via CREB:CBP-dependent transcriptional activation. *J. Neurosci.* **27**, 6128–6140 (2007).
52. E. K. Schmidt, G. Clavarino, M. Ceppi, P. Pierre, SUnSET, a nonradioactive method to monitor protein synthesis. *Nat. Methods* **6**, 275–277 (2009).
53. E. Santini, T. N. Huynh, A. F. MacAskill, A. G. Carter, P. Pierre, D. Ruggero, H. Kaphzan, E. Klann, Exaggerated translation causes synaptic and behavioural aberrations associated with autism. *Nature* **493**, 411–415 (2013).
54. C. A. Hoeffer, W. Tang, H. Wong, A. Santillan, R. J. Patterson, L. A. Martinez, M. V. Tejada-Simon, R. Paylor, S. L. Hamilton, E. Klann, Removal of FKBP12 enhances mTOR-raptor interactions, LTP, memory, and perseverative/repetitive behavior. *Neuron* **60**, 832–845 (2008).
55. A. M. M. Oliveira, J. D. Hawk, T. Abel, R. Havekes, Post-training reversible inactivation of the hippocampus enhances novel object recognition memory. *Learn. Mem.* **17**, 155–160 (2010).
56. R. Havekes, D. A. Canton, A. J. Park, T. Huang, T. Nie, J. P. Day, L. A. Guercio, Q. Grimes, V. Luczak, I. H. Gelman, G. S. Baillie, J. D. Scott, T. Abel, Gravin orchestrates protein kinase A and β 2-adrenergic receptor signaling critical for synaptic plasticity and memory. *J. Neurosci.* **32**, 18137–18149 (2012).

Acknowledgments: We would like to thank N. Grissom, T. Jongens, and N. Sonenberg for editorial input on the manuscript; P. Hernandez for technical support; and C. Hoeffer and E. Klann for fruitful discussions and advice. **Funding:** This work was supported by NIH grants F32MH099730 (to J.C.T.), R21MH102703 (to T.A.), P01AG017628 (to T.A.), K12GM081259 (to J.C.T.; principal investigator, Y. Paterson), and T32NS007413 (to L.P.; principal investigator, M. Robinson). This work was also supported by the Kwanjeong Educational Foundation (to A.J.P.) and the Brush Family Professorship (to T.A.). **Author contributions:** J.C.T., L.P., M.E.W., A.J.P., and R.H. designed and performed the sleep deprivation studies. J.C.T., E.J.D., E.v.T., and C.W.C. performed surgeries, along with the biochemistry and behavioral experiments. J.C.T. and S.G.P. performed the qPCR experiments. J.C.T., E.G., and P.P. designed the in vivo translation experiment. J.C.T. and J.H. performed the statistical analyses. J.C.T. and T.A. wrote the paper. **Competing interests:** The authors declare that they have no competing interests. **Data and materials availability:** The pAAV₉-CaMKII α 0.4-eIF4EBP2-HA virus is available by contacting T.A.

Submitted 22 September 2015

Accepted 6 April 2016

Final Publication 26 April 2016

10.1126/scisignal.aad4949

Citation: J. C. Tudor, E. J. Davis, L. Peixoto, M. E. Wimmer, E. van Tilborg, A. J. Park, S. G. Poplawski, C. W. Chung, R. Havekes, J. Huang, E. Gatti, P. Pierre, T. Abel, Sleep deprivation impairs memory by attenuating mTORC1-dependent protein synthesis. *Sci. Signal.* **9**, ra41 (2016).

Sleep deprivation impairs memory by attenuating mTORC1-dependent protein synthesis

Jennifer C. Tudor, Emily J. Davis, Lucia Peixoto, Mathieu E. Wimmer, Erik van Tilborg, Alan J. Park, Shane G. Poplawski, Caroline W. Chung, Robbert Havekes, Jiayan Huang, Evelina Gatti, Philippe Pierre and Ted Abel

Sci. Signal. **9** (425), ra41.
DOI: 10.1126/scisignal.aad4949

How sleep deprivation impairs memory

Sleep deprivation impairs learning and memory. Tudor *et al.* (see also the Focus by Sweatt and Hawkins) found that sleep deprivation in mice suppressed activation of the kinase complex mTORC1 and consequently protein synthesis in hippocampal neurons, which impaired their memory. Restoring protein synthesis by increasing the amount of phosphorylated 4EBP2 protein in the hippocampus—a function normally performed by mTORC1—protected mice from the memory impairment caused by sleep deprivation. The findings reveal a molecular mechanism by which sleep loss impairs memory.

ARTICLE TOOLS

<http://stke.sciencemag.org/content/9/425/ra41>

SUPPLEMENTARY MATERIALS

<http://stke.sciencemag.org/content/suppl/2016/04/22/9.425.ra41.DC1>

RELATED CONTENT

<http://stke.sciencemag.org/content/sigtrans/9/425/fs7.full>
<http://stke.sciencemag.org/content/sigtrans/8/382/ra61.full>
<http://stke.sciencemag.org/content/sigtrans/7/355/re12.full>
<http://science.sciencemag.org/content/sci/350/6259/379.full>
<http://science.sciencemag.org/content/sci/350/6256/82.full>
<http://stm.sciencemag.org/content/scitransmed/7/291/291ec96.full>
<http://stke.sciencemag.org/content/sigtrans/9/426/ec107.abstract>
<http://science.sciencemag.org/content/sci/352/6285/517.full>
<http://stke.sciencemag.org/content/sigtrans/9/430/fs10.full>
<http://stke.sciencemag.org/content/sigtrans/9/430/ra57.full>
<http://stke.sciencemag.org/content/sigtrans/9/434/ec153.abstract>
<http://stke.sciencemag.org/content/sigtrans/9/441/ec188.abstract>
<http://science.sciencemag.org/content/sci/353/6300/687.full>
<http://stke.sciencemag.org/content/sigtrans/10/460/eaam6552.full>
<http://stke.sciencemag.org/content/sigtrans/10/463/eaai9219.full>
<http://science.sciencemag.org/content/sci/355/6324/507.full>
<http://science.sciencemag.org/content/sci/355/6324/457.full>
<http://science.sciencemag.org/content/sci/355/6324/511.full>
<http://stke.sciencemag.org/content/sigtrans/10/465/eaam9084.full>
<http://stke.sciencemag.org/content/sigtrans/10/504/eaan0665.full>
<http://stke.sciencemag.org/content/sigtrans/10/505/eaao2327.full>
<http://stke.sciencemag.org/content/sigtrans/10/505/eaar4559.full>
<http://science.sciencemag.org/content/sci/351/6269/aaa5694.full>
<http://science.sciencemag.org/content/sci/352/6285/550.full>
<http://stke.sciencemag.org/content/sigtrans/11/532/eaau2601.full>

Use of this article is subject to the [Terms of Service](#)

Science Signaling (ISSN 1937-9145) is published by the American Association for the Advancement of Science, 1200 New York Avenue NW, Washington, DC 20005. The title *Science Signaling* is a registered trademark of AAAS.

Copyright © 2016, American Association for the Advancement of Science

REFERENCES

This article cites 56 articles, 15 of which you can access for free
<http://stke.sciencemag.org/content/9/425/ra41#BIBL>

PERMISSIONS

<http://www.sciencemag.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of Service](#)

Science Signaling (ISSN 1937-9145) is published by the American Association for the Advancement of Science, 1200 New York Avenue NW, Washington, DC 20005. The title *Science Signaling* is a registered trademark of AAAS.

Copyright © 2016, American Association for the Advancement of Science